



Always ask your healthcare professional which product is right for your child.

## Dosing Information for Healthcare Professionals

*Infants'*  
**TYLENOL**

*Children's*  
**TYLENOL**

Use this chart to determine the proper dose of **TYLENOL**® for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 4 hours as needed.  
Do not give more than 5 doses in 24 hours.**

WEIGHT	AGE	Infants' <b>TYLENOL</b> ® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL)	Children's <b>TYLENOL</b> ® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL or 1 tsp)
6-11 lbs	0-3 mos	1.25 mL	—
12-17 lbs	4-11 mos	2.5 mL	—
18-23 lbs	12-23 mos	3.75 mL	—
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoonful

**Remind parents and caregivers to:**

- Read and follow the label on all **TYLENOL**® products
- Take every 4 hours as needed.
- Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device that comes with a specific product

**All Infants' **TYLENOL**® and Children's **TYLENOL**® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)**

## Dosing Information for Healthcare Professionals

*Children's*  
**Motrin**

*Concentrated*  
**Motrin**  
*Infants' Drops*

Use this chart to determine the proper dose of **MOTRIN**® for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 6-8 hours as needed.  
Do not give more than 4 doses in 24 hours.**

WEIGHT	AGE	Concentrated <b>MOTRIN</b> ® Infants' Drops Active Ingredient: Ibuprofen 50 mg (NSAID)* (in each 1.25 mL)	Children's <b>MOTRIN</b> ® Oral Suspension Active Ingredient: Ibuprofen 100 mg (NSAID)* (in each 5 mL or 1 tsp)
6-11 lbs	0-5 mos	—	—
12-17 lbs	6-11 mos	1.25 mL	—
18-23 lbs	12-23 mos	1.875 mL	—
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoonful

**Remind parents and caregivers to:**

- Read and follow the label on all **MOTRIN**® products
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID\* (prescription or non-prescription)  
— **MOTRIN**® contains ibuprofen
- Use only the dosing device that comes with a specific product

\*Nonsteroidal anti-inflammatory drug  
©McNEIL-PPC, Inc. 2013 • CTCM-0144